

OIP
MAY 17 2002
PATENT TRADEMARK OFFICE

Page 1 of 2

Atty. Docket No.: 3130-001 Application No.: 09/933,610

Applicant: Zhong, et al.

Filing Date: August 21, 2001

Group Art Unit: 1732

U.S. PATENT DOCUMENTS


Examiner Initial*	Document Number	Date	Name	Class	Sub Class	Filing Date If Appropriate

FOREIGN PATENT DOCUMENTS

Document Number	Date	Country	Class	Sub Class	Translation Yes or No

OTHER DOCUMENTS (Including Author, Title, Date, Pertinent Pages, Etc.)

SA	De Smet et al., <i>Characterization of Intracellular Inclusions Formed by Pseudomonas oleovorans During Growth on Octane</i> , <u>Journal of Bacteriology</u> , p. 870-878 (May 1983) /
	Wallen et al., <i>Poly-β-hydroxyalkanoate from Activated Sludge</i> , <u>Environmental Science & Technology</u> , Vol. 8, pp. 576-579, (June 1974) /
	Steinbuchel et al., <i>A Pseudomonas strain accumulating polyesters of 3-hydroxybutyric acid and medium-chain-length 3-hydroxyalkanoic acids</i> <u>Applied Microbiology and Biotechnology</u> , pp. 691-697 (1992) /
	Valentin et al., <i>Identification of 4-hydroxyvaleric acid as a constituent of biosynthetic polyhydroxyalkanoic acids from bacteria</i> , <u>Applied Microbiology and Biotechnology</u> , pp. 507-514, (1992) /
	Valentin et al., <i>Identification of 4-hydroxyhexanoic acid as a new constituent of biosynthetic polyhydroxyalkanoic acids from bacteria</i> , <u>Applied Microbiology and Biotechnology</u> , pp. 710-716, (1994) /
	Lee et al., <i>Biosynthesis of copolyesters consisting of 3-hydroxybutyric acid and medium-chain-length 3-hydroxyalkanoic acids from 1,3-butanediol or from 3-hydroxybutyrate by Pseudomonas sp. A33</i> , <u>Applied Microbiology and Biotechnology</u> , pp. 901-909, (1995) /
	Kato et al., <i>Production of a novel copolyester of 3-hydroxybutyric acid and medium-chain-length 3-hydroxyalkanoic acids by Pseudomonas sp. 61-3 from sugars</i> , <u>Applied Microbiology and Biotechnology</u> , pp. 363-370, (1996) /
	Valentin et al., <i>Identification of 5-hydroxyhexanoic acid, 4-hydroxyheptanoic acid and 4-hydroxyoctanoic acid as new constituents of bacterial polyhydroxyalkanoic acids</i> , <u>Applied Microbiology and Biotechnology</u> , pp. 261-267, (1996) /
	Brandl et al., <i>Ability of the phototrophic bacterium Rhodospirillum rubrum to produce various poly (β-hydroxyalkanoates): potential sources for biodegradable polyesters</i> , <u>Int. J. Biol. Macromol.</u> , Vol. 11, pp. 49-55, (1989) /
	Amos et al., <i>Composition of poly-β-hydroxyalkanoate from Syntrophomonas wolfei grown on unsaturated fatty acid substrates</i> , <u>Archives of Microbiology</u> , pp. 103-105, (1991) /
	Byrom, D. <i>Miscellaneous Biomaterials</i> , ICI BioProducts and Fine Chemicals, pp. 335-359. /
SA	Hocking et al., <u>Biopolyesters</u> , pp. 48-96. /



SA	Lafferty et al., <u>Microbial Production of Poly-β-hydroxybutyric Acid</u> , Chapter 6, pp. 137-176. /		
	Gross et al., <u>Polymerization of β-Monosubstituted-β-propiolactones Using Trialkylaluminum-Water Catalytic Systems and Polymer Characterization</u> , <u>Macromolecules</u> , pp. 2657-2668, (1988) /		
	Dubois et al., <u>Macromolecular Engineering of Polylactones and Polylactides</u> , <u>Macromolecules</u> , pp. 4407-4412, (1993) /		
	Borgne et al., <u>Stereoelective polymerization of β-butyrolactone</u> , <u>Polymer</u> , Vol. 30, pp. 2312-2319, (December 1989) /		
	Tanahashi et al., <u>Thermal Properties and Stereoregularity of Poly(3-hydroxybutyrate) Prepared from Optically Active β-Butyrolactone with a Zinc-Based Catalyst</u> , <u>Macromolecules</u> , Vol. 24, pp. 5732-5733 (1991) /		
	Hori et al., <u>Ring-Opening Copolymerization of Optically Active β-Butyrolactone with Several Lactones Catalyzed by Distannoxane Complexes: Synthesis of New Biodegradable Polyesters</u> , <u>Macromolecules</u> , Vol. 26, pp. 4388-4390 / (1993)		
	Kemnitzer, <u>Preparation of Predominantly Syndiotactic Poly (β-hydroxybutyrate) by the Tributyltin Methoxide Catalyzed Ring-Opening Polymerization of Racemic β-Butyrolactone</u> , <u>Macromolecules</u> , Vol. 26, pp. 1221-1229, (1993) /		
	Hori et al., <u>Ring-Opening Polymerization of Optically Active β-Butyrolactone Using Distannoxane Catalysts: Synthesis of High Molecular Weight Poly(3-hydroxybutyrate)</u> , <u>Macromolecules</u> , Vol. 26, pp. 5533-5534 (1993) /		
	Hocking et al., <u>Syndiotactic poly[(R,S)-β-hydroxybutyrate] isolated from methylaluminoxane-catalyzed polymerization</u> , <u>Polymer Bulletin</u> , Vol. 30, pp. 163-170 (1993) /		
	Xie et al., <u>Ring-Opening Polymerization of β-Butyrolactone by Thermophilic Lipases</u> , <u>Macromolecules</u> , Vol. 30, pp. 6997-6998 (1997) /		
	Jesudason et al., <u>Synthetic Poly[(R,S)-β-hydroxyalkanoates] with Butyl and Hexyl Side Chains</u> , <u>Macromolecules</u> , Vol. 27, pp. 2595-2602 (1994) /		
	<u>Reports on Progress in Polymer Physics in Japan</u> , Vol. 37, pp.128-129 (1994) /		
	Kharas et al., <u>Polymers of Lactic Acid</u> 94-137 (No date available) /		
	Abe et al., <u>Biosynthesis from gluconate of a random copolyester consisting of 3-hydroxy-butyrate and medium-chain-length 3-hydroxyalkanoates by Pseudomonas sp. 61-3</u> , <u>Int. J. Biol. Macromol.</u> , Vol. 16, pp. 36, 38, 40, (1994) /		
SA	Muller et al., <u>Poly(hydroxyalkanoates): A Fifth Class of Physiologically Important Organic Biopolymers?</u> , <u>Angew. Chem. Int. Ed. Engl.</u> , Vol. 32, pp. 477-502 (1993)		
Examiner	Sam ACQUAH	Date Considered	4/20/03
*Examiner:	Initial if reference considered, whether or not citation is in conformance with MPEP 609; draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.		
Form PTO 1449		Patent and Trademark Office - U.S. Department of Commerce	